


# MATH 2205: Calculus II

---

Libao Jin


June 20, 2019

University of Wyoming



# Review for Midterm Exam 2


---



# Review for Midterm Exam 2

---

## Integration Techniques



## Proposition (Basic Integration Formulas)

$$(a) \int k dx = kx + C, k \in \mathbb{R} \text{ (} k \text{ is real).}$$

$$(b) \int x^p dx = \frac{x^{p+1}}{p+1} + C, p \neq -1 \in \mathbb{R}.$$

$$(c) \int e^{ax} dx = \frac{1}{a} e^{ax} + C.$$

$$(d) \int \frac{1}{x} dx = \ln |x| + C.$$

$$(e) \int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a} + C.$$

$$(f) \int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} + C.$$

$$(g) \int \frac{1}{x\sqrt{x^2 - a^2}} dx = \frac{1}{a} \sec^{-1} \left| \frac{x}{a} \right| + C, a > 0.$$



### Proposition (Basic Integration Formulas (continued))

$$(a) \int \cos ax \, dx = \frac{1}{a} \sin ax + C.$$

$$(b) \int \sin ax \, dx = -\frac{1}{a} \cos ax + C.$$

$$(c) \int \sec^2 ax \, dx = \frac{1}{a} \tan ax + C.$$

$$(d) \int \csc^2 ax \, dx = -\frac{1}{a} \cot ax + C.$$

$$(e) \int \sec ax \tan ax \, dx = \frac{1}{a} \sec ax + C.$$

$$(f) \int \csc ax \cot ax \, dx = -\frac{1}{a} \csc ax + C.$$

# Integration by Parts

## Theorem (Integration by Parts)

Suppose that  $u$  and  $v$  are differentiable functions. Then

$$\int u \, dv = uv - \int v \, du.$$

## Theorem (Integration by Parts for Definite Integrals)

Let  $u$  and  $v$  be differentiable. Then

$$\int_a^b u(x)v'(x) \, dx = u(x)v(x) \Big|_a^b - \int_a^b v(x)u'(x) \, dx.$$

## Powers of $\sin x$ or $\cos x$

### Procedure

Strategies for evaluating integrals of the form  $\int \sin^m x \, dx$  or  $\int \cos^n x \, dx$ , where  $m$  and  $n$  are positive integers, using trigonometric identities.

- (a) Integrals involving odd powers of  $\cos x$  (or  $\sin x$ ) are most easily evaluated by splitting off a single factor of  $\cos x$  (or  $\sin x$ ). For example, rewrite  $\cos^5 x$  as  $\cos^4 x \cdot \cos x$ .
- (b) With even positive powers of  $\sin x$  or  $\cos x$ , we use the half-angle formulas

$$\sin^2 \theta = \frac{1 - \cos 2\theta}{2} \quad \text{and} \quad \cos^2 \theta = \frac{1 + \cos 2\theta}{2},$$

to reduce the powers in the integrand.



## Products of Powers of $\sin x$ and $\cos x$

### Procedure

Strategies for evaluating integrals of the form  $\int \sin^m x \cos^n x dx$ .

- (a) When  $m$  is odd and positive,  $n$  real. Split off  $\sin x$ , rewrite the resulting even power of  $\sin x$  in terms of  $\cos x$ , and then use  $u = \cos x$ .
- (b) When  $n$  is odd and positive,  $m$  real. Split off  $\cos x$ , rewrite the resulting even power of  $\cos x$  in terms of  $\sin x$ , and then use  $u = \sin x$ .
- (c) When  $m, n$  are both even and nonnegative. Use half-angle formulas to transform the integrand into polynomial in  $\cos 2x$  and apply the preceding strategies once again to powers of  $\cos 2x$  greater than 1.



## Proposition (Reduction Formulas)

Assume  $n$  is a positive integer.

$$(a) \int \sin^n x \, dx = -\frac{\sin^{n-1} x \cos x}{n} + \frac{n-1}{n} \int \sin^{n-2} x \, dx.$$

$$(b) \int \cos^n x \, dx = \frac{\cos^{n-1} x \sin x}{n} + \frac{n-1}{n} \int \cos^{n-2} x \, dx.$$

$$(c) \int \tan^n x \, dx = \frac{\tan^{n-1} x}{n-1} - \int \tan^{n-2} x \, dx, n \neq 1.$$

$$(d) \int \sec^n x \, dx = \frac{\sec^{n-2} x \tan x}{n-1} + \frac{n-2}{n-1} \int \sec^{n-2} x \, dx, n \neq 1.$$

# Trigonometric Substitutions

## Proposition (Integral contains $a^2 - x^2$ )

Let  $x = a \sin \theta$ ,  $-\pi/2 \leq \theta \leq \pi/2$  for  $|x| \leq a$ . Then

$$a^2 - x^2 = a^2 - a^2 \sin^2 \theta = a^2(1 - \cos^2 \theta) = a^2 \cos^2 \theta.$$

## Procedure (Partial Fractions with Simple Linear Factors)

Suppose  $f(x) = p(x)/q(x)$ , where  $p$  and  $q$  are polynomials with no common factors and with the degree of  $p$  less than the degree of  $q$ . Assume that  $q$  is the product of simple linear factors. The partial fraction decomposition is obtained as follows.

- Factor the denominator  $q$*  in the form  $(x-r_1)(x-r_2)\cdots(x-r_n)$ , where  $r_1, \dots, r_n$  are real numbers.
- Partial fraction decomposition.* Form the partial fraction decomposition by writing

$$\frac{p(x)}{q(x)} = \frac{A_1}{(x-r_1)} + \frac{A_2}{(x-r_2)} + \cdots + \frac{A_n}{(x-r_n)}.$$



## Procedure (Partial Fractions with Simple Linear Factors (continued))

- (c) *Clear denominators.* Multiply both sides of the equation in Step (b) by  $q(x) = (x - r_1)(x - r_2) \cdots (x - r_n)$ , which produces conditions for  $A_1, \dots, A_n$ .
- (d) *Solve for coefficients.* Equate like powers of  $x$  in Step (c) to solve for the undetermined coefficients  $A_1, \dots, A_n$ .



## Procedure (Partial Fractions for Repeated Linear Factors)

Suppose the repeated linear factor  $(x - r)^m$  appears in the denominator of a proper rational function in reduced form. The partial fraction decomposition has a partial fraction for each power of  $(x - r)$  up to and including the  $m$ th power; that is, the partial fraction decomposition contains the sum

$$\frac{A_1}{(x - r)} + \frac{A_2}{(x - r)^2} + \cdots + \frac{A_m}{(x - r)^m},$$

where  $A_1, \dots, A_m$  are constants to be determined.



# Partial Fractions

## Procedure (Partial Fractions with Simple Irreducible Quadratic Factors)

Suppose a simple irreducible factor  $ax^2 + bx + c$  appears in the denominator of a proper rational function in reduced form. The partial fraction decomposition contains a term of the form

$$\frac{Ax + B}{ax^2 + bx + c},$$

where  $A$  and  $B$  are unknown coefficients to be determined.

## Proposition

The quadratic polynomial  $ax^2 + bx + c$  is irreducible if and only if its discriminant is negative, i.e.,

$$\Delta = b^2 - 4ac < 0.$$



## Proposition (Partial Fraction Decomposition)

Let  $f(x) = p(x)/q(x)$  be a proper rational function in reduced form. Assume the denominator  $q$  has been factored completely over the real numbers and  $m$  is a positive integer.

- (a) *Simple linear factor.* A factor  $x - r$  in the denominator requires the partial fraction  $\frac{A}{x - r}$ .
- (b) *Repeated linear factor.* A factor  $(x - r)^m$  with  $m > 1$  in the denominator requires the partial fractions

$$\frac{A_1}{(x - r)} + \frac{A_2}{(x - r)^2} + \frac{A_3}{(x - r)^3} + \cdots + \frac{A_m}{(x - r)^m}.$$



## Proposition (Partial Fraction Decomposition (continued))

(c) *Simple irreducible quadratic factor.* An irreducible factor  $ax^2 + bx + c$  in the denominator requires the partial fraction

$$\frac{Ax + B}{ax^2 + bx + c}.$$

(d) *Repeated irreducible quadratic factor.* An irreducible factor  $(ax^2 + bx + c)^m$  with  $m > 1$  in the denominator requires the partial fractions

$$\frac{A_1x + B_1}{ax^2 + bx + c} + \frac{A_2x + B_2}{(ax^2 + bx + c)^2} + \cdots + \frac{A_mx + B_m}{(ax^2 + bx + c)^m}.$$





## Definition (Absolute and Relative Error)

Suppose  $c$  is a computed numerical solution to a problem having an exact solution  $x$ . There are two common measures of the error in  $c$  as an approximation to  $x$ :

$$\text{absolute error} = |c - x|$$

and

$$\text{relative error} = \frac{c - x}{x}, (\text{if } x \neq 0).$$



## Definition (Midpoint Rule)

Suppose  $f$  is defined an integrable on  $[a, b]$ . The *Midpoint Rule approximation* to  $\int_a^b f(x) dx$  using  $n$  equally spaced subintervals on  $[a, b]$  is

$$\begin{aligned}M(n) &= f(m_1)\Delta x + f(m_2)\Delta x + \cdots + f(m_n)\Delta x \\ &= \sum_{k=1}^n f\left(\frac{x_{k-1} + x_k}{2}\right) \Delta x,\end{aligned}$$

where  $\Delta x = (b - a)/n$ ,  $x_0 = a$ ,  $x_k = a + k\Delta x$ , and  $m_k = (x_{k-1} + x_k)/2 = a + (k - 1/2)\Delta x$  is the midpoint of  $[x_{k-1}, x_k]$ , for  $k = 1, \dots, n$ .



## Definition (Trapezoid Rule)

Suppose  $f$  is defined and integrable on  $[a, b]$ . The *Trapezoid Rule approximation* to  $\int_a^b f(x) dx$  using  $n$  equally spaced subintervals on  $[a, b]$  is

$$T(n) = \left[ \frac{1}{2}f(x_0) + \sum_{k=1}^{n-1} f(x_k) + \frac{1}{2}f(x_n) \right] \Delta x.$$

where  $\Delta x = (b - a)/n$  and  $x_k = a + k\Delta x$ , for  $k = 0, 1, 2, \dots, n$ .



## Definition (Simpson's Rule)

Suppose  $f$  is defined and integrable on  $[a, b]$  and  $n \geq 2$  is an even integer. The *Simpson's Rule approximation* to  $\int_a^b f(x) dx$  using  $n$  equally spaced subintervals on  $[a, b]$  is

$$S(n) = \sum_{k=0}^{n/2-1} [f(x_{2k}) + 4f(x_{2k+1}) + f(x_{2k+2})] \frac{\Delta x}{3}.$$

where  $n$  is an even integer,  $\Delta x = (b - a)/n$ , and  $x_k = a + k\Delta x$ , for  $k = 0, 1, \dots, n$ .

## Definition (Improper Integrals over Infinite Intervals)

(a) If  $f$  is continuous on  $[a, \infty)$ , then

$$\int_a^{\infty} f(x) dx = \lim_{b \rightarrow \infty} \int_a^b f(x) dx.$$

(b) If  $f$  is continuous on  $(-\infty, b]$ , then

$$\int_{-\infty}^b f(x) dx = \lim_{a \rightarrow -\infty} \int_a^b f(x) dx.$$



## Definition (Improper Integrals over Infinite Intervals (continued))

(c) If  $f$  is continuous on  $(-\infty, \infty)$ , then

$$\int_{-\infty}^{\infty} f(x) dx = \lim_{a \rightarrow -\infty} \int_a^c f(x) dx + \lim_{b \rightarrow \infty} \int_c^b f(x) dx,$$

where  $c$  is any real number.

If the limits in the above cases exist, then the improper integrals *converge*; otherwise, they *diverge*.



## Definition (Improper Integrals with an Unbounded Integrand)

(a) Suppose  $f$  is continuous on  $(a, b]$  with  $\lim_{x \rightarrow a^+} f(x) = \pm\infty$ . Then

$$\int_a^b f(x) dx = \lim_{c \rightarrow a^+} \int_c^b f(x) dx.$$

(b) Suppose  $f$  is continuous on  $[a, b)$  with  $\lim_{x \rightarrow b^-} f(x) = \pm\infty$ . Then

$$\int_a^b f(x) dx = \lim_{c \rightarrow b^-} \int_a^c f(x) dx.$$



## Definition (Improper Integrals with an Unbounded Integrand (continued))

(c) Suppose  $f$  is continuous on  $[a, b]$  except at the interior point  $p$  where  $f$  is unbounded. Then

$$\int_a^b f(x) dx = \lim_{c \rightarrow p^-} \int_a^c f(x) dx + \lim_{d \rightarrow p^+} \int_d^b f(x) dx.$$

If the limits in above cases exist, then the improper integrals *converge*; otherwise, they *diverge*.



## Definition

The *order* of a differential equation is the highest order appearing on a derivative in the equation. For example, the equations  $y' + 4y = \cos x$  and  $y' = 0.1y(100 - y)$  are first order, and  $y'' + 16y = 0$  is second order.

## Definition (Linear Differential Equations)

The first-order *linear* differential equations have the form

$$y'(x) + p(x)y(x) = f(x),$$

and the second-order *linear* differential equations have the form

$$y''(x) + p(x)y'(x) + q(x)y(x) = f(x),$$

where  $p$ ,  $q$ , and  $f$  are given functions that depend only on the independent variable  $x$ .



## Definition

A differential equation is often accompanied by *initial conditions* that specify the values of  $y$ , and possibly its derivatives, at a particular point. In general, an  $n$ th-order equation requires  $n$  initial conditions.



## Definition

A differential equation, together with the appropriate number of initial conditions, is called an *initial value problem*. A typical first-order initial value problem has the form

$$y'(t) = F(t, y)$$

Differential equation

$$y(0) = A$$

Initial condition

where  $A$  is given and  $F$  is a given expression that involves  $t$  and/or  $y$ ,

## Proposition (Solution of a First-Order Linear Differential Equation)

The general solution of the first-order equation  $y'(t) = ky + b$ , where  $k$  and  $b$  are specified real numbers, is  $y = Ce^{kt} - b/k$ , where  $C$  is an arbitrary constant. Given an initial condition, the value of  $C$  may be determined.



## Definition (Separable First-Order Differential Equations)

If the first-order differential equation can be written in the form  $g(y)y'(t) = h(t)$ , in which the terms that involve  $y$  appear on one side of the equation *separated* from the terms that involve  $t$ , is said to be *separable*. We can solve the equation by integrating both sides of the equation with respect to  $t$ :


$$\int g(y)y'(t) dt = \int h(t) dt \implies \int g(y) dy = \int h(t) dt.$$



# Review for Midterm Exam 2

---

## Sequences and Infinite Series



## Definition (Sequence)

A *sequence*  $\{a_n\}$  is an ordered list of numbers of the form

$$\{a_1, a_2, a_3, \dots, a_n, \dots\}.$$

- (a) A sequence may be generated by a *recurrence relation* of the form  $a_{n+1} = f(a_n)$ , for  $n = 1, 2, 3, \dots$ , where  $a_1$  is given.
- (b) A sequence may also be defined with an *explicit formula* of the form  $a_n = f(n)$ , for  $n = 1, 2, 3, \dots$





# Limit of a Sequence

## Definition (Limit of a Sequence)

If the terms of a sequence  $\{a_n\}$  approach a unique number  $L$  as  $n$  increases – that is, if  $a_n$  can be made arbitrarily close to  $L$  by taking  $n$  sufficiently large – then we say  $\lim_{n \rightarrow \infty} a_n = L$  exists, and the sequence *converges* to  $L$ . If the terms of the sequence do not approach a single number as  $n$  increases, the sequence has no limit, and the sequence *diverges*.

## Theorem (Limits of Sequences from Limits of Functions)

Suppose  $f$  is a function such that  $f(n) = a_n$  for all positive integers  $n$ . If  $\lim_{x \rightarrow \infty} f(x) = L$ , then the limit of the sequence  $\{a_n\}$  is also  $L$ .



## Theorem (Limits of Linear Functions)

*Let  $a$ ,  $b$ , and  $m$  be real numbers. For linear functions*

$$f(x) = mx + b,$$

$$\lim_{x \rightarrow a} f(x) = f(a) = ma + b.$$



## Theorem (Limit Laws)

Assume  $\lim_{x \rightarrow a} f(x)$  and  $\lim_{x \rightarrow a} g(x)$  exist. The following properties hold, where  $c$  is a real number, and  $m > 0$  and  $n > 0$  are integers.

(a) *Sum:*  $\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x).$

(b) *Difference:*  $\lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x).$

(c) *Constant multiple:*  $\lim_{x \rightarrow a} cf(x) = c \lim_{x \rightarrow a} f(x).$

(d) *Product:*  $\lim_{x \rightarrow a} [f(x)g(x)] = \left[ \lim_{x \rightarrow a} f(x) \right] \left[ \lim_{x \rightarrow a} g(x) \right].$

(e) *Quotient:*  $\lim_{x \rightarrow a} \left[ \frac{f(x)}{g(x)} \right] = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)},$  provided  $\lim_{x \rightarrow a} g(x) \neq 0.$

(f) *Power:*  $\lim_{x \rightarrow a} [f(x)]^n = \left[ \lim_{x \rightarrow a} f(x) \right]^n.$



## Theorem (Limits of Polynomial and Rational Functions)

Assume  $p$  and  $q$  are polynomials and  $a$  is a constant.

(a) Polynomial functions:  $\lim_{x \rightarrow a} p(x) = p(a)$ .

(b) Rational functions:  $\lim_{x \rightarrow a} \frac{p(x)}{q(x)} = \frac{p(a)}{q(a)}$ , provided  $q(a) \neq 0$ .



## Theorem (The Squeeze Theorem)

Assume the function  $f$ ,  $g$ , and  $h$  satisfy  $f(x) \leq g(x) \leq h(x)$  for all values of  $x$  near  $a$ , except possibly at  $a$ . If

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L, \text{ then } \lim_{x \rightarrow a} g(x) = L.$$



## Theorem (Limits at Infinity of Powers and Polynomials)

Let  $n$  be a positive integer and let  $p$  be the polynomial

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_2 x^2 + a_1 x + a_0, \text{ where } a_n \neq 0.$$

(a)  $\lim_{x \rightarrow \pm\infty} x^n = \infty$  when  $n$  is even.

(b)  $\lim_{x \rightarrow \infty} x^n = \infty$  and  $\lim_{x \rightarrow -\infty} x^n = -\infty$  when  $n$  is odd.

(c)  $\lim_{x \rightarrow \pm\infty} \frac{1}{x^n} = \lim_{x \rightarrow \pm\infty} x^{-n} = 0$ .

(d)  $\lim_{x \rightarrow \pm\infty} p(x) = \lim_{x \rightarrow \pm\infty} a_n x^n = \pm\infty$ , depending on the degree of the polynomial and the sign of the leading coefficient  $a_n$ .



## Theorem (End Behavior of $e^x$ , $e^{-x}$ , and $\ln x$ )

The end behavior for  $e^x$  and  $e^{-x}$  on  $(-\infty, \infty)$  and  $\ln x$  on  $(0, \infty)$  is given by the following limits (see Figure 1):

$$\lim_{x \rightarrow \infty} e^x = \infty$$

$$\lim_{x \rightarrow -\infty} e^x = 0$$

$$\lim_{x \rightarrow \infty} e^{-x} = 0$$


$$\lim_{x \rightarrow -\infty} e^{-x} = \infty$$

$$\lim_{x \rightarrow 0^+} \ln x = -\infty$$

$$\lim_{x \rightarrow \infty} \ln x = \infty.$$



# Limits of Functions



**Figure 1:** Graphs of  $e^x$ ,  $e^{-x}$ ,  $\ln x$ :  $y = e^{-x}$  and  $y = e^x$  are symmetric about  $y$ -axis, and  $y = e^x$  and  $y = \ln x$  are symmetric about  $y = x$ .



## Theorem (L'Hôpital's Rule)

Suppose  $f$  and  $g$  are differentiable on an open interval  $I$  containing  $a$  with  $g'(x) \neq 0$  on  $I$  when  $x \neq a$ .

(a) If  $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$ , then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)},$$

provided the limit on the right exists (or is  $\pm\infty$ ). The rule also applies if  $x \rightarrow a$  is replaced with  $x \rightarrow \pm\infty$ ,  $x \rightarrow a^+$ ,  $x \rightarrow a^-$ .

(b) If  $\lim_{x \rightarrow a} f(x) = \pm\infty$  and  $\lim_{x \rightarrow a} g(x) = \pm\infty$ , then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)},$$

provided the limit on the right exists (or is  $\pm\infty$ ). The rule also applies if  $x \rightarrow a$  is replaced with  $x \rightarrow \pm\infty$ ,  $x \rightarrow a^+$ ,  $x \rightarrow a^-$ .

## Theorem (Limit Laws for Sequences)

Assume that the sequences  $\{a_n\}$  and  $\{b_n\}$  have limits  $A$  and  $B$ , respectively. Then

$$(a) \lim_{n \rightarrow \infty} (a_n \pm b_n) = A \pm B.$$

$$(b) \lim_{n \rightarrow \infty} ca_n = cA, \text{ where } c \text{ is a real number.}$$

$$(c) \lim_{n \rightarrow \infty} a_n b_n = AB.$$

$$(d) \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{A}{B} \text{ provided } B \neq 0.$$

## Definition

- (a)  $\{a_n\}$  is *increasing* if  $a_{n+1} > a_n$ ; for example,  $\{0, 1, 2, 3, \dots\}$ .
- (b)  $\{a_n\}$  is *nondecreasing* if  $a_{n+1} \geq a_n$ ; for example,  $\{0, 1, 1, 1, 2, 2, 3, \dots\}$ .
- (c)  $\{a_n\}$  is *decreasing* if  $a_{n+1} < a_n$ ; for example,  $\{2, 1, 0, -2, \dots\}$ .
- (d)  $\{a_n\}$  is *nonincreasing* if  $a_{n+1} \leq a_n$ ; for example,  $\{2, 1, 1, 0, -2, -2, -3, \dots\}$ .
- (e)  $\{a_n\}$  is *monotonic* if it is either nonincreasing or nondecreasing (it moves in one direction).
- (f)  $\{a_n\}$  is *bounded* if there is number  $M$  such that  $|a_n| \leq M$ , for all relevant values of  $n$ .



# Squeeze Theorem for Sequences

## Theorem (Squeeze Theorem for Sequences)

Let  $\{a_n\}$ ,  $\{b_n\}$ , and  $\{c_n\}$  be sequences with  $a_n \leq b_n \leq c_n$  for all integers  $n$  greater than some index  $N$ . If  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = L$ , then  $\lim_{n \rightarrow \infty} b_n = L$ .

## Theorem (Bounded Monotonic Sequences)

A bounded monotonic sequence converges.



## Theorem (Growth Rates of Sequences)

The following sequences are ordered according to increasing growth rates as  $n \rightarrow \infty$ ; that is, if  $\{a_n\}$  appears before  $\{b_n\}$  in the list, then  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0$  and  $\lim_{n \rightarrow \infty} \frac{b_n}{a_n} = \infty$ :

$$\{\ln^q n\} \ll \{n^p\} \ll \{n^p \ln^r n\} \ll \{n^{p+s}\} \ll \{b^n\} \ll \{n!\} \ll \{n^n\}.$$

The ordering applies for positive real numbers  $p, q, r, s$  and  $b > 1$ .



## Definition (Infinite series)

Given a sequence  $\{a_1, a_2, a_3, \dots\}$ , the sum of its terms

$$a_1 + a_2 + a_3 + \cdots = \sum_{k=1}^{\infty} a_k$$

is called an *infinite series*.



# Sequence of Partial Sums

## Definition (Sequence of Partial Sums)

The *sequence of partial sums*  $\{S_n\}$  associated with this series has the terms

$$S_1 = a_1$$

$$S_2 = a_1 + a_2$$

$$S_3 = a_1 + a_2 + a_3$$

$\vdots$

$$S_n = a_1 + a_2 + a_3 + \cdots + a_n = \sum_{k=1}^n a_k, \text{ for } n = 1, 2, 3, \dots$$



# Sequence of Partial Sums and Infinite Series

## Proposition

If the sequence of partial sums  $\{S_n\}$  has a limit  $L$ , the infinite series *converges* to that limit, and we write

$$\sum_{k=1}^{\infty} a_k = \lim_{n \rightarrow \infty} \sum_{k=1}^n a_k = \lim_{n \rightarrow \infty} S_n = L.$$

If the sequence of partial sums diverges, the infinite series also *diverges*.





# Geometric Sequences

## Definition (Geometric Sequences)

A sequence has the form  $\{r^n\}$  or  $\{ar^n\}$ , where the ratio  $r$ ,  $a$  are real numbers, is called a geometric sequence.

## Theorem (Geometric Sequences)

Let  $r$  be a real number. Then

$$\lim_{n \rightarrow \infty} r^n = \begin{cases} 0 & \text{if } |r| < 1, \\ 1 & \text{if } r = 1, \\ \text{does not exist} & \text{if } r \leq -1 \text{ or } r > 1. \end{cases}$$

If  $r > 0$ , then  $\{r^n\}$  is a monotonic sequence. If  $r < 0$ , then  $\{r^n\}$  oscillates.



## Theorem (Geometric Series)

Let  $a \neq 0$  and  $r$  be real numbers. If  $|r| < 1$ , then  $\sum_{k=0}^{\infty} ar^k = \frac{a}{1-r}$ .

If  $|r| \geq 1$ , then the series diverges. More generally,


$$\sum_{k=m}^{\infty} ar^k = \frac{ar^m}{1-r}.$$



# Review for Midterm Exam 2

---

Algebra



# Exponents and Radicals

$$(a) \frac{1}{x^a} = x^{-a}.$$

$$(b) \sqrt[n]{x} = x^{1/n}.$$

$$(c) x^{a+b} = x^a x^b.$$


$$(d) x^{a-b} = \frac{x^a}{x^b}.$$

$$(e) x^{ab} = (x^a)^b.$$

$$(f) x^{m/n} = \sqrt[n]{x^m} = (\sqrt[n]{x})^m.$$

$$(g) (xy)^a = x^a y^a.$$

$$(h) \left(\frac{x}{y}\right)^a = \frac{x^a}{y^a}.$$



# Logarithm

$$(a) y = a^x \implies x = \log_a y.$$

$$(b) \log_e x = \ln x.$$

$$(c) \log_b(xy) = \log_b x + \log_b y.$$

$$(d) \log_b \frac{x}{y} = \log_b x - \log_b y.$$

$$(e) \log_b(x^p) = p \log_b x.$$

$$(f) \log_b(x^{1/p}) = \frac{1}{p} \log_b x.$$

$$(g) \log_b x = \frac{\log_k x}{\log_k b}.$$



## Factoring Formulas

$$(a) \quad a^2 - b^2 = (a - b)(a + b).$$

$$(b) \quad a^3 - b^3 = (a - b)(a^2 + ab + b^2).$$


$$(c) \quad a^n - b^n = (a - b)(a^{n-1} + a^{n-2}b + a^{n-3}b^2 + \dots + ab^{n-2} + b^{n-1}).$$



# Binomials

$$(a) \quad (a \pm b)^2 = a^2 \pm 2ab + b^2.$$

$$(b) \quad (a \pm b)^3 = a^3 \pm 3a^2b + 3ab^2 \pm b^3.$$



## Completing the Square: $x^2 \pm bx + c$

Given that  $(x \pm p)^2 = x^2 \pm 2px + p^2$ .

$$\begin{aligned}x^2 \pm bx + c &= x^2 \pm 2\frac{b}{2}x + c \\&= x^2 \pm 2\frac{b}{2}x + \left(\frac{b}{2}\right)^2 + c - \left(\frac{b}{2}\right)^2 \\&= \left(x \pm \frac{b}{2}\right)^2 + c - \frac{b^2}{4}.\end{aligned}$$





## Completing the Square: $ax^2 \pm bx + c$

$$\begin{aligned}ax^2 \pm bx + c &= a \left( x^2 \pm \frac{b}{a}x \right) + c \\&= a \left( x^2 \pm 2 \frac{b}{2a}x \right) + c \\&= a \left[ x^2 \pm 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 \right] + c - a \left( \frac{b}{2a} \right)^2 \\&= a \left( x \pm \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a} \\&= (\sqrt{a})^2 \left( x \pm \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a} \\&= \left( \sqrt{a}x \pm \frac{\sqrt{ab}}{2a} \right)^2 + c - \frac{b^2}{4a} \\&= \left( \sqrt{a}x \pm \frac{b}{2\sqrt{a}} \right)^2 + c - \frac{b^2}{4a}.\end{aligned}$$



# Quadratic Formula

The solutions of  $ax^2 + bx + c = 0$  are

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

