


MATH 3341: Introduction to Scientific Computing Lab

Libao Jin

University of Wyoming

November 04, 2020





Lab 11: MATLAB Integration Routines &
Gauss Quadrature





Built-in Integration Routines



polyint: Indefinite Integral

- $I = \text{polyint}(p, c)$: indefinite integral of polynomial p with c being the constant.
- $I = \text{polyint}(p)$: same as $\text{polyint}(p, 0)$.
- Example: $\int 3x^2 + 2x + 1 dx = x^3 + x^2 + x + C$.

```
p = [3,2,1]
```

```
I1 = polyint(p, 1) % [1,1,1,1]
```

```
I2 = polyint(p, 2) % [1,1,1,2]
```

```
I3 = polyint(p) % [1,1,1,0]
```

```
polyval(I1, 0) % 1
```

```
polyval(I2, 0) % 2
```

```
polyval(I3, 0) % 0
```



polyint: Definite Integral

- Fundamental Theorem of Calculus (FTOC):

$$\int_a^b p'(x) dx = p(x) \Big|_a^b = p(b) - p(a).$$

- Example: $\int_0^2 3x^2 + 2x + 1 dx = x^3 + x^2 + x + C \Big|_{x=0}^{x=2} = 14.$

`p = [3,2,1]`

`P = polyint(p)` % [1,1,1,0]

`I = polyval(P, 2) - polyval(P, 0)` % 14



trapz: Trapezoidal numerical integration

- $I = \text{trapz}(x, y)$ computes the integral of y with respect to x using the trapezoidal method, x and y must be vectors of the same length.
- Let $X = [x_1, x_2]$, $Y = [y_1, y_2]$, it is actually a trapezoid, where y_1 and y_2 are the lengths for the bases and $x_2 - x_1$ is the height. Then

$$I = \frac{(x_2 - x_1)(y_1 + y_2)}{2}.$$

- Let $X = [x_1, x_2, \dots, x_n]$, $Y = [y_1, y_2, \dots, y_n]$, then

$$I = \sum_{i=1}^{n-1} \frac{(x_{i+1} - x_i)(y_{i+1} + y_i)}{2} = \frac{1}{2} \sum_{i=1}^{n-1} (x_{i+1} - x_i)(y_{i+1} + y_i).$$



cumtrapz: Cumulative trapezoidal numerical integration

- $I = \text{cumtrapz}(x, y)$ computes the cumulative integral of y with respect to x using trapezoidal integration.
- Example:

```
x = [1,2,3];
```

```
y = [1,2,3];
```

```
I1 = cumtrapz(x, y) % [0, 1.5000, 4.0000]
```

```
I2 = [trapz([1], [1]),  
      trapz([1,2], [1,2]),  
      trapz([1,2,3], [1,2,3])] % [0, 1.5000, 4.0000]
```



Numerically evaluate integral - 1D

- `I = integral(f, a, b)` approximates the integral of function `f` from `a` to `b` using global adaptive quadrature and default error tolerances. `f` must be a function handle, `a` and `b` can be `-Inf` or `Inf`.

- Example: $\int_0^2 3x^2 + 2x + 1 dx = x^3 + x^2 + x + C \Big|_{x=0}^{x=2} = 14.$

```
f = @(x) 3 * x.^2 + 2 * x + 1;
```

```
a = 0;
```

```
b = 2;
```

```
I = integral(f, a, b)           % 14.0000
```



Numerically evaluate integral - 2D

- `I = integral2(f,xmin,xmax,ymin,ymax)` approximates the integral of $f(x,y)$ over the planar region $x_{\min} \leq x \leq x_{\max}$ and $y_{\min}(x) \leq y \leq y_{\max}(x)$. f is a function handle, y_{\min} and y_{\max} may each be a scalar value or a function handle.

- Example:

$$\int_0^2 \int_0^x 6y + 2 \, dy \, dx = \int_0^2 3x^2 + 2x \, dx = x^3 + x^2 + C \Big|_{x=0}^{x=2} = 12.$$

```
f = @(x, y) 6 * y + 2;
```

```
xmin = 0;
```

```
xmax = 2;
```

```
ymin = 0;
```

```
ymin = @(x) x;
```

```
I = integral2(f, xmin, xmax, ymin, ymax); % 12.0000
```

Numerically evaluate integral - 3D

- `I = integral3(f, xmin, xmax, ymin, ymax, zmin, zmax)` approximates the integral of $f(x,y,z)$ over the region $x_{\min} \leq x \leq x_{\max}$, $y_{\min}(x) \leq y \leq y_{\max}(x)$, and $z_{\min}(x,y) \leq z \leq z_{\max}(x,y)$. f is a function handle, y_{\min} , y_{\max} , z_{\min} , and z_{\max} may each be a scalar value or a function handle.

- Example:
$$\int_0^2 \int_0^x \int_{-2y+2}^{4y+4} 1 \, dz \, dy \, dx = \int_0^2 \int_0^x 6y + 2 \, dy \, dx =$$

$$\int_0^2 3x^2 + 2x \, dx = x^3 + x^2 + C \Big|_{x=0}^{x=2} = 12.$$

```
f = @(x, y, z) ones(size(z));
```

```
xmin = 0; xmax = 2;
```

```
ymin = 0; ymax = @(x) x;
```

```
zmin = @(x,y) -2 * y + 4;
```

```
zmax = @(x,y) 4 * y + 4;
```

```
I = integral3(f, xmin, xmax, ymin, ymax, zmin, zmax);
```





Gauss-Legendre Quadrature



Gauss-Legendre Quadrature on $[-1, 1]$

Integration of $f(x)$ on the interval $[-1, 1]$ using Gauss Quadrature is given by

$$\int_{-1}^1 f(x) dx \approx \sum_{i=1}^n w_i f(x_i),$$

where w_i and x_i are chosen so the integration rule is exact for the largest class of polynomials. $f(x)$ is well-approximated by polynomial on $[-1, 1]$, the associated orthogonal polynomials are *Legendre polynomial*, denoted by $P_n(x)$. With the n -th polynomial normalized to give $P_n(1) = 1$, the i -th Gauss node, x_i , is the i -th root of P_n and the weights are given by the formula (Abramowitz & Stegun 1972, p. 887):

$$w_i = \frac{2}{(1 - x_i^2)[P_n'(x_i)]^2}.$$



Gauss-Legendre Quadrature on $[a, b]$

To approximate the integral on the general interval $[a, b]$, we need to use the change of variables as follows:

$$\frac{t-a}{b-a} = \frac{x-(-1)}{1-(-1)} = \frac{x+1}{2} \implies t = \frac{b-a}{2}x + \frac{b+a}{2}, -1 \leq x \leq 1$$
$$\implies dt = \frac{b-a}{2}dx.$$

So the Gauss Quadrature on a general interval $[a, b]$ is given by

$$\int_a^b f(t) dt = \int_{-1}^1 f\left(\frac{b-a}{2}x + \frac{b+a}{2}\right) \frac{b-a}{2} dx$$
$$\approx \sum_{i=1}^n w_i f\left(\frac{b-a}{2}x_i + \frac{b+a}{2}\right) \frac{b-a}{2}.$$



Gauss-Legendre Quadrature on $[a, b]$

Let

$$g(x) = f\left(\frac{b-a}{2}x + \frac{b+a}{2}\right) \frac{b-a}{2},$$

then

$$\int_a^b f(t) dt = \int_{-1}^1 g(x) dx \approx \sum_{i=1}^n w_i g(x_i).$$

