


MATH 3341: Introduction to Scientific Computing Lab

Libao Jin

University of Wyoming

March 24, 2021





Lab 09: Ill-conditioned Matrices and Finite
Precision Arithmetic



The background features a large, faint watermark of the University of Wyoming seal. The seal is circular with a rope-like border. Inside the border, the words "UNIVERSITY OF WYOMING" are written in an arc at the top, and "1886" is at the bottom. In the center, there is an open book with a quill pen resting on it, and the word "EQUALITY" is written below the book.

Ill-Conditioned Matrices



Vector Norm

Let $\mathbf{x} = [x_1, x_2, \dots, x_n]^T \in \mathbb{R}^n$.

- $\text{norm}(\mathbf{x}, 1)$: $\|\mathbf{x}\|_1 = \sum_{i=1}^n |x_i|$.
- $\text{norm}(\mathbf{x}, 2)$: $\|\mathbf{x}\|_2 = \left(\sum_{i=1}^n x_i^2 \right)^{1/2} = (\mathbf{x} \cdot \mathbf{x})^{1/2}$.
- $\text{norm}(\mathbf{x}, \text{inf})$: $\|\mathbf{x}\|_\infty = \max_{i=1, \dots, n} \{|x_i|\}$.



Matrix Norm

Let $\mathbf{x} = [x_1, x_2, \dots, x_n]^T \in \mathbb{R}^n$ and $A \in \mathbb{R}^{m \times n}$.

- $\text{norm}(A, 1)$: $\|A\|_1 = \max_{\substack{\forall \mathbf{x} \in \mathbb{R}^n \\ \mathbf{x} \neq \mathbf{0}}} \frac{\|A\mathbf{x}\|_1}{\|\mathbf{x}\|_1} = \max_{1 \leq j \leq n} \sum_{i=1}^m |a_{ij}|.$

- $\text{norm}(A, 2)$:

$$\|A\|_2 = \max_{\substack{\forall \mathbf{x} \in \mathbb{R}^n \\ \mathbf{x} \neq \mathbf{0}}} \frac{\|A\mathbf{x}\|_2}{\|\mathbf{x}\|_2} = \sigma_{\max}(A) = \sqrt{\lambda_{\max}(A^T A)},$$

$\sigma_{\max}(A)/\lambda_{\max}(A)$ means the largest singular value/eigenvalue of matrix A .

- $\text{norm}(A, \text{inf})$: $\|A\|_{\infty} = \max_{\substack{\forall \mathbf{x} \in \mathbb{R}^n \\ \mathbf{x} \neq \mathbf{0}}} \frac{\|A\mathbf{x}\|_{\infty}}{\|\mathbf{x}\|_{\infty}} = \max_{i=1, \dots, m} \sum_{j=1}^n |a_{ij}|.$

- $\text{norm}(A, \text{'fro'})$: $\|A\|_F = \left(\sum_{i=1}^n \sum_{j=1}^n a_{ij}^2 \right)^{1/2}$, Frobenius norm.



Condition Number

The *condition number* of nonsingular matrix A relative to the norm $\|\cdot\|$ is

$$\kappa(A) = \|A\| \cdot \|A^{-1}\|.$$

- If the condition number is high, then the matrix is said to be *ill-conditioned*.
- If $\kappa(A) = \infty$, then the matrix A is singular, i.e., the matrix is not invertible.



cond: condition number with respect to inversion

- `cond(A, 1)`: 1-norm condition number of A.
- `cond(A, 2)`: 2-norm condition number of A, i.e.,
$$\kappa_2(A) = \|A\|_2 \|A^{-1}\|_2 = \frac{\sigma_{\max}(A)}{\sigma_{\min}(A)}.$$
- `cond(A, inf)`: Infinity-norm condition number of A.
- `cond(A, 'fro')`: Frobenius-norm condition number of A.
- `cond(A)`: same as `cond(A, 2)`.
- Example

```
A = magic(5);  
condA1 = cond(A, 2)  
condA2 = norm(A, 2) * norm(inv(A), 2)  
condA3 = max(sqrt(eig(A'*A))) * max(sqrt(eig(inv(A'*A))))  
condA4 = max(sqrt(eig(A'*A))) / min(sqrt(eig(A'*A)))
```



Ill-Conditioned Matrix: Hilbert Matrix

A Hilbert matrix is a square matrix with elements defined by

$$H_{ij} = \frac{1}{i + j - 1}.$$

For example, a 3×3 Hilbert matrix is

$$H_{3 \times 3} = \begin{bmatrix} 1 & 1/2 & 1/3 \\ 1/2 & 1/3 & 1/4 \\ 1/3 & 1/4 & 1/5 \end{bmatrix}.$$

Note that this matrix is symmetric and positive definite.



hilb: Hilbert matrix and invhilb: inverse Hilbert matrix

- `hilb(n)`: the n -by- n matrix with elements $1/(i+j-1)$, which is a famous example of a ill-conditioned matrix.
- `invhilb(n)`: the inverse of the n -by- n Hilbert matrix. The result is exact for n less than about 15.
- Example:

```
H = hilb(10);  
invH1 = inv(H);  
invH2 = invhilb(10);  
norm(invH1 - invH2)
```



Ill-Conditioned Linear System

A linear system $Ax = b$ is said to be ill-conditioned if A is a ill-conditioned matrix. The typical numerical methods for solving linear systems such as Jacobi method, Gauss-Seidel method would become unreliable. Example: $Hx = b \implies x = H^{-1}b$.

```
n = 10;  
H = hilb(n);  
invH = invhilb(n);  
b = rand(n, 1);  
x = invH * b;  
x1 = inv(H) * b;  
x2 = H \ b;  
norm(x - x1)  
norm(x - x2)
```



The background features a large, faint watermark of the University of Wyoming seal. The seal is circular with a rope-like border. Inside the border, the words "UNIVERSITY OF WYOMING" are written in an arc at the top, and "1886" is at the bottom. In the center, there is an open book with a quill pen resting on it, and the word "EQUALITY" is written below the book.

Finite Precision Arithmetic



Finite Precision Arithmetic

Computers can only store values up to a certain level of accuracy. Past this level, the computer will round values, thus causes the round-off error. What this means is that arithmetic does not work exactly as we expect. Namely, arithmetic is no longer commutative, associative, or distributive. The lab exercises will demonstrate some of the issues that arise.



IEEE 754

<https://babbage.cs.qc.cuny.edu/IEEE-754/>

